

# WHEELCHAIR CPR

A method of preparing a cardiac arrest casualty in a wheelchair for CPR

**DANGER**  
Remove hazards

**RESPONSE**  
Try to get a response

**SEND FOR HELP**  
Ambulance & assistance

**AIRWAY**  
Open with head tilt/chin lift

**BREATHING**  
Look, listen and feel

**MAKE ROOM BEHIND THE CHAIR - APPLY THE BRAKES - REMOVE ACCESSORIES**  
(armrests, anti-tip devices, food tray etc).

## One first aider

If possible, secure the casualty to the chair  
(ie around the chest and chair back with a belt)  
Stand behind the chair  
Hold on to the handles

## Two first aiders

One on each side of the chair  
Kneel down on one knee  
(with the foot at the rear of the chair on the ground)  
Place your arm under the casualty's armpit and hold onto the handle  
The other hand supports the casualty's head

## More than two first aiders

If several people can help:  
Make room on the floor  
Carefully lift the casualty from the wheelchair  
Place on his/her back on the ground

**LOWER THE CHAIR BACKWARDS TO THE GROUND**

**REMOVE FROM THE CHAIR**

Support under armpits, pull/slide out of the chair onto the ground

**START CPR**

**Ensure Safety for Self and Others**

**Dial Triple Zero (000) for an Ambulance**

## Signs & Symptoms

Cannot be woken.  
No purposeful response.  
Not breathing normally or at all.

There may be methods other than these.  
Use good manual handling techniques.  
Do not attempt if beyond your strength capabilities.  
Avoiding injury to the casualty and the first aider is the priority.  
Try not to bang the casualty's head.

Only do CPR in the chair if it has a hard back.

*May not be suitable for motorised wheelchairs.*

*Do not do this in a plastic chair.*